arXiv:math/0609749v1 [math.NT] 27 Sep 2006

Asymptotics of abelian group-partitions and associated Dirichlet series

Tetsuya MOMOTANI

February 2, 2008

Abstract

We introduce a notion of a group-partition for a finite Abelian group, which is a generalized notion of the standard partition. To obtain asymptotic distributions of group-partition, we study the Dirichlet series for group-partitions by employing the generating function of the plane partition. [1]

1 Introduction

Let $a(n)$ be the number of isomorphic classes of Abelian groups of order n It is known (Cf. [Sa]) that

$$\sum_{\substack{n\le x \\ n\equiv k \pmod{j}}} a(n) = C(j)x + o(x) \qquad \text{as } x\to\infty. \tag{1.1}$$

Here j and k are positive integers such that $(j,k)=1$ and $C(j)$ is a positive constant.

The aim of this paper is a generalization of (1.1) by introducing a new notion which we call a group partition; for an isomorphic class G of finite Abelian groups, a sequence (G_j) of isomorphic classes of finite Abelian groups is called a group-partition of G if (G_j) satisfies

$$G = G_1 \oplus G_2 \oplus G_3 \oplus \cdots \oplus G_r, \quad G_1 \supseteq G_2 \supseteq G_3 \supseteq \cdots \supseteq G_r.$$

Let $\pi_r(G)$ be the number of group-partitions of G with at most r factors. Let $a_r(n)$ be an arithmetic function defined by $a_r(n) := \sum_{|G|=n} \pi_r(G)$ where

---

[1] 2000 *Mathematics Subject Classification* : Primary 11N37, 20K01.

Key words: partition, plane partition, finite Abelian groups, zeta functions.

G runs over isomorphic classes of finite Abelian groups of order n. Note that $a_1(n) = a(n)$. Then the main result claims that

$$\sum_{\substack{n\le x \\ n\equiv k \ (\bmod\ j)}} a_r(n) = C_r(j)x + o(x) \qquad \text{as } x \to \infty. \tag{1.2}$$

Here $C_r(j)$ is a positive constant explicitly expressed in terms of the Riemann zeta function and the Euler function (Theorem 4.3).

To prove (1.2), we introduce the Dirichlet series defined by

$$J_r(s,\chi) := \sum_{n=1}^{\infty} \chi(n)a_r(n)n^{-s} \qquad \mathrm{Re}(s) > 1,$$

where χ is a Dirichlet character. Using plane partitions and its generating functions, we show that $a_r(n)$ is multiplicative in the sence of $a_r(mn) = a_r(m)a_r(n)$ when $(m,n) = 1$. From this function, we see that $J_r(s,\chi)$ has an Euler product expression and an analytic continuation to $\mathrm{Re}(s) > 0$. Then the asymptotic distribution (1.2) follows from the Ikehara-Wiener theorem by the standard technique.

2 Group-partitions and plane partitions

Let n be a positive integer and (n_j) be a sequence of non-negative integers such that

$$n = n_1 + n_2 + n_3 + \cdots, \quad n_1 \ge n_2 \ge n_3 \ge \cdots.$$

Then (n_j) is called a partition of n, and we write $(n_j) \vdash n$. We denote by $P(n)$ the number of all partitions of n.

As a generalization of notion of partitions for positive integers, we define group-partitions for finite Abelian groups.

Definition 2.1. *Let G be an isomorphic class of finite Abelian groups. Suppose that (G_j) is a sequence of isomorphic classes of finite Abelian groups such that*

$$G = G_1 \oplus G_2 \oplus G_3 \oplus \cdots, \quad G_1 \supseteq G_2 \supseteq G_3 \supseteq \cdots.$$

Then we call (G_j) a group-partition of G, and write $(G_j) \vdash G$. For a positive integer r, we define $\pi_r(G)$ as the number of group-partitions of G with at most r factors. We also define $\pi_\infty(G)$ as the number of all group-partitions of G. □

We give some examples.

Example 1. Suppose that $G = (\mathbb{Z}/p\mathbb{Z})^n$ with a prime number p and a positive integer n. We put $G_j = (\mathbb{Z}/p\mathbb{Z})^{n_j}$. Then $(G_j) \vdash G \Leftrightarrow (n_j) \vdash n$. Hence we have $\pi_\infty(G) = P(n)$. The decomposition theorem of finite Abelian groups states that every finite Abelian group can be expressed as the direct sum of $(\mathbb{Z}/p^m\mathbb{Z})^n$'s. Hence the group-partition is a generalization of the usual partition.

Example 2. If $G = (\mathbb{Z}/p\mathbb{Z})^2 \oplus (\mathbb{Z}/p^2\mathbb{Z})$ with a prime number p, then $\pi_1(G) = 1$, $\pi_2(G) = 2$, $\pi_3(G) = \pi_4(G) = \cdots = \pi_\infty(G) = 3$. In fact, all of group-partitions of G are given as follows.

(1) $G = G_1$, where $G_1 = (\mathbb{Z}/p\mathbb{Z})^2 \oplus (\mathbb{Z}/p^2\mathbb{Z})$.

(2) $G = G_1 \oplus G_2$, $\quad G_1 = (\mathbb{Z}/p\mathbb{Z}) \oplus (\mathbb{Z}/p^2\mathbb{Z})$, $G_2 = (\mathbb{Z}/p\mathbb{Z})$.

(3) $G = G_1 \oplus G_2 \oplus G_3$, $\quad G_1 = (\mathbb{Z}/p^2\mathbb{Z})$, $G_2 = (\mathbb{Z}/p\mathbb{Z})$, $G_3 = (\mathbb{Z}/p\mathbb{Z})$.

Example 3. If $G = (\mathbb{Z}/p\mathbb{Z})^3 \oplus (\mathbb{Z}/q\mathbb{Z})^2$ with prime numbers $p \neq q$, then $\pi_1(G) = 1$, $\pi_2(G) = 4$, $\pi_3(G) = \pi_4(G) = \cdots = \pi_\infty(G) = 6$.

(1) $G = G_1$, where $G_1 = (\mathbb{Z}/p\mathbb{Z})^3 \oplus (\mathbb{Z}/q\mathbb{Z})^2$.

(2) $G = G_1 \oplus G_2$, $\quad G_1 = (\mathbb{Z}/p\mathbb{Z})^3 \oplus (\mathbb{Z}/q\mathbb{Z})$, $G_2 = (\mathbb{Z}/q\mathbb{Z})$.

(3) $G = G_1 \oplus G_2$, $\quad G_1 = (\mathbb{Z}/p\mathbb{Z})^2 \oplus (\mathbb{Z}/q\mathbb{Z})^2$, $G_2 = (\mathbb{Z}/p\mathbb{Z})$,

(4) $G = G_1 \oplus G_2$, $\quad G_1 = (\mathbb{Z}/p\mathbb{Z})^2 \oplus (\mathbb{Z}/q\mathbb{Z})$, $G_2 = (\mathbb{Z}/p\mathbb{Z}) \oplus (\mathbb{Z}/q\mathbb{Z})$,

(5) $G = G_1 \oplus G_2 \oplus G_3$, $G_1 = (\mathbb{Z}/p\mathbb{Z}) \oplus (\mathbb{Z}/q\mathbb{Z})^2$, $G_2 = (\mathbb{Z}/p\mathbb{Z})$, $G_3 = (\mathbb{Z}/p\mathbb{Z})$.

(6) $G = G_1 \oplus G_2 \oplus G_3$, $G_1 = (\mathbb{Z}/p\mathbb{Z}) \oplus (\mathbb{Z}/q\mathbb{Z})$, $G_2 = (\mathbb{Z}/p\mathbb{Z}) \oplus (\mathbb{Z}/q\mathbb{Z})$, $G_3 = (\mathbb{Z}/p\mathbb{Z})$.

In general, it is easy to see the following.

Lemma 2.2. *Suppose that $p_1, p_2, p_3, \cdots$ are distinct prime numbers. If G_{p_j} is an Abelian p_j-group, then we have $\pi_r(G_{p_1} \oplus G_{p_2} \oplus G_{p_3} \oplus \cdots) = \pi_r(G_{p_1}) \times \pi_r(G_{p_2}) \times \pi_r(G_{p_3}) \times \cdots$.* □

Remark. In the recent work [AS], a subgroup tower of Abelian groups is introduced and studied. A subgroup tower of a group G is defined as a pair of groups $(G_1, G_2, G_3, \cdots)$ where $G_1 = G$ and $G_1 \supseteq G_2 \supseteq G_3 \supseteq \cdots$. We note that definitions of a subgroup tower and a group-partition are similar but different.

Let

$$a_r(n) := \sum_{|G|=n} \pi_r(G),$$

where G runs over isomorphic classes of Abelian groups of order n. We note that $a_1(n) = a(n)$ since $\pi_1(G) = 1$ for any G. In order to investigate this function, we recall several results about the plane partition. Let n be a positive integer and (λ_{jk}) be a matrix where λ_{jk} are non-negative integers such that

$$\sum_{j\geq 0, k\geq 0} \lambda_{jk} = n, \quad \lambda_{j1} \geq \lambda_{j2} \geq \lambda_{j3} \geq \cdots, \quad \lambda_{1k} \geq \lambda_{2k} \geq \lambda_{3k} \geq \cdots.$$

Then (λ_{jk}) is called a plane partition of n. Let $\mathrm{PL}_r(n)$ denote the number of plane partitions of n with at most r-rows. Let $\mathrm{PL}_\infty(n)$ denote the number of all plane partitions of n. It is known (Cf. [An] p.184) that generating functions of the r-rowed plane partition $\mathrm{PL}_r(n)$ and the plane partition $\mathrm{PL}_\infty(n)$ are given by

(2.1)

$$\sum_{n=0}^{\infty} \mathrm{PL}_r(n) q^n = \prod_{m=1}^{\infty} (1-q^m)^{-\min\{m,r\}}, \qquad \sum_{n=0}^{\infty} \mathrm{PL}_\infty(n) q^n = \prod_{m=1}^{\infty} (1-q^m)^{-m},$$

for $|q| < 1$. Now we show the following proposition which plays a key role in this paper.

Proposition 2.3. *Let p be a prime number and n be a positive integer. For $r = 1, 2, 3, \cdots$, or $r = \infty$, we have*

$$a_r(p^n) = \sum_{|G|=p^n} \pi_r(G) = \mathrm{PL}_r(n). \tag{2.2}$$

Proof. We show that there is a one-to-one correspondence between group-partitions of Abelian p-groups with order p^n and plane partitions of n. First we suppose that (λ_{jk}) is a r-rowed plane partition of n, that is

$$\sum_{j=1}^{r} \sum_{k=1}^{n} \lambda_{jk} = n, \quad \lambda_{j1} \geq \lambda_{j2} \geq \cdots \geq \lambda_{jn} \geq 0, \quad \lambda_{1k} \geq \lambda_{2k} \geq \cdots \geq \lambda_{rk} \geq 0.$$

We put

$$G_j := (\mathbb{Z}/p\mathbb{Z})^{\lambda_{j1}-\lambda_{j2}} \oplus (\mathbb{Z}/p^2\mathbb{Z})^{\lambda_{j2}-\lambda_{j3}} \oplus \cdots \oplus (\mathbb{Z}/p^n\mathbb{Z})^{\lambda_{jn}},$$

and $G := G_1 \oplus G_2 \oplus \cdots \oplus G_r$. Then G_j, G are Abelian p-groups such that $|G| = p^n$ and $(G_j) \vdash G$. Actually, the condition $\lambda_{1k} \geq \lambda_{2k} \geq \cdots \geq \lambda_{rk}$ means that $G_1 \supseteq G_2 \supseteq G_3 \supseteq \cdots \supseteq G_r$.

Conversely, we assume that G is a Abelian group of order p^n, and (G_j) is a group-partition of G with at most r factors. Let μ_{jk} by

$$G_j = (\mathbb{Z}/p\mathbb{Z})^{\mu_{j1}} \oplus (\mathbb{Z}/p^2\mathbb{Z})^{\mu_{j2}} \oplus \cdots \oplus (\mathbb{Z}/p^n\mathbb{Z})^{\mu_{jn}},$$

and $\lambda_{jk} := \sum_{m=k}^{n} \mu_{jm}$. Then we see that $\lambda_{j1} \geq \lambda_{j2} \geq \cdots \geq \lambda_{jn} \geq 0$. Since we have $|G| = p^n$, it follows that $\sum_{j=1}^{r} \sum_{k=1}^{n} \lambda_{jk} = n$. Moreover $G_1 \supseteq G_2 \supseteq G_3 \supseteq \cdots \supseteq G_r$ means that $\lambda_{1k} \geq \lambda_{2k} \geq \cdots \geq \lambda_{rk}$. Hence (λ_{jk}) is a plane partition of n with at most r rows. This completes the proof of the proposition. ☐

Corollary 2.4. *The function $a_r(n)$ is multiplicative.*

Proof. Let $n = p_1^{m_1} p_2^{m_2} p_3^{m_3} \cdots$. Using Lemma 2.2 and (2.2), we have

$$\begin{aligned} a_r(n) &= \sum_{|G|=p_1^{m_1} p_2^{m_2} p_3^{m_3} \cdots} \pi_r(G) = \sum_{|G_1|=p_1^{m_1}} \sum_{|G_2|=p_2^{m_2}} \cdots \pi_r(G_1 \oplus G_2 \oplus \cdots) \\ &= \Big(\sum_{|G_1|=p_1^{m_1}} \pi_r(G_1) \Big) \times \Big(\sum_{|G_2|=p_2^{m_2}} \pi_r(G_2) \Big) \times \cdots = \mathrm{PL}_r(m_1) \times \mathrm{PL}_r(m_2) \times \cdots \\ &= a_r(p_1^{m_1}) \cdot a_r(p_2^{m_2}) \cdot a_r(p_3^{m_3}) \cdot \cdots . \end{aligned}$$

This shows that $a_r(n)$ is multiplicative. ☐

Corollary 2.5. *For any $\epsilon > 0$, we have*

$$a_r(n) \leq a_\infty(n) = O_\epsilon(n^\epsilon). \tag{2.3}$$

Proof. It is known (Cf. [An] p.199) that

$$\mathrm{PL}_\infty(n) \sim (\zeta(3)^7 \cdot 2^{-11})^{1/36} \cdot n^{-25/36} \cdot \exp\{3 \cdot 2^{-2/3} \cdot \zeta(3)^{1/3} \cdot n^{2/3} + 2c\}$$

as $n \to \infty$. Here

$$c = \int_0^\infty \frac{y \log y}{e^{2\pi y} - 1} dy$$

and $\zeta(s)$ is the Riemann zeta function. Using this, we estimate roughly as $\mathrm{PL}_\infty(n) \leq \exp\{Cn^{2/3}\}$ where C is a positive number. Then for $\epsilon > 0$ and $n = p_1^{m_1} p_2^{m_2} p_3^{m_3} \cdots$, we have

$$\begin{aligned} 0 \leq \frac{a_\infty(n)}{n^\epsilon} &= \prod_{p_i \leq e^{C/\epsilon}} \frac{\mathrm{PL}_\infty(m_i)}{p_i^{m_i \epsilon}} \times \prod_{p_i > e^{C/\epsilon}} \frac{\mathrm{PL}_\infty(m_i)}{p_i^{m_i \epsilon}} \\ &\leq \prod_{p_i \leq e^{C/\epsilon}} \frac{\exp(C m_i^{2/3})}{p_i^{m_i \epsilon}} \times \prod_{p_i > e^{C/\epsilon}} \frac{\exp(C m_i^{2/3})}{\exp(C m_i)} \\ &\leq \prod_{p_i \leq e^{C/\epsilon}} \frac{\exp(C m_i^{2/3})}{p_i^{m_i \epsilon}} \leq D_\epsilon. \end{aligned}$$

Here $D_\epsilon > 0$ depends only on ϵ. Hence the result follows. ☐

3 Dirichlet series and Euler products

We first recall the classical Dirichlet L-function. Let χ be a Dirichlet character modulo j and $L(s,\chi) := \sum_{n=1}^{\infty} \chi(n)n^{-s}$ be the Dirichlet L-function. Then $L(s,\chi)$ is defined for $\mathrm{Re}(s) > 1$, and has the Euler product expression: $L(s,\chi) = \prod_p (1-\chi(p)p^{-s})^{-1}$. If χ_0 is the principal character modulo j, then the Dirichlet L-function $L(s,\chi_0)$ is essentially the Riemann zeta function:

$$L(s,\chi_0) = \prod_{p|j}(1-p^{-s}) \times \zeta(s).$$

For a non-principal character $\chi \neq \chi_0$, it is known that the Drichlet L-function $L(s,\chi)$ can be analytically continued as an entire function.

We next consider the Dirichlet series:

$$J_r(s,\chi) := \sum_{n=1}^{\infty} \chi(n)a_r(n)n^{-s} \qquad (r = 1,2,3,\cdots,\infty).$$

We see from (2.3) that this series $J_r(s,\chi)$ converges absolutely for $\mathrm{Re}(s) > 1$. Further, we show that $J_r(s,\chi)$ has the Euler product expression as follows.

Proposition 3.1. *For $r = 1,2,3,\cdots$, or $r = \infty$, the function $J_r(s,\chi)$ has the Euler product expression:*

(3.1)

$$J_r(s,\chi) = \prod_{m=1}^{\infty} \prod_{p:\ \text{prime}} (1-\chi(p)^m p^{-ms})^{-\min\{m,r\}} = \prod_{m=1}^{\infty} L(ms,\chi^m)^{\min\{m,r\}},$$

$$J_\infty(s,\chi) = \prod_{m=1}^{\infty} \prod_{p:\ \text{prime}} (1-\chi(p)^m p^{-ms})^{-m} = \prod_{m=1}^{\infty} L(ms,\chi^m)^m.$$

Here the product converges absolutely for $\mathrm{Re}(s) > 1$.

Proof. Since the sum $\sum_{m=1}^{\infty}\sum_p mp^{-m\sigma}$ converges for $\sigma > 1$, it follows that the Euler product of (3.1) converges absolutely for $\mathrm{Re}(s) > 1$. For a fixed prime number p, we see from (2.1) and (2.2) that

$$\prod_{m=1}^{\infty}(1-\chi(p)^m p^{-ms})^{-\min\{m,r\}} = \sum_{n=0}^{\infty} \mathrm{PL}_r(n)\chi(p)^n p^{-ns} = \sum_{n=0}^{\infty} a_r(p^n)\chi(p^n)p^{-ns}.$$

Since $a_r(n)$ is a multiplicative function, we obtain (3.1). □

Corollary 3.2. *The function $J_r(s,\chi)$ defined for* $\mathrm{Re}(s) > 1$ *has a meromorphic continuation to the half plane* $\mathrm{Re}(s) > 0$.

Proof. Using Proposition 3.1, we have

(3.2)

$$J_r(s,\chi) = \prod_{m=1}^{M-1} L(ms,\chi^m)^{\min\{m,r\}} \times \prod_{m=M}^{\infty} \prod_{p} (1-\chi(p)^m p^{-ms})^{-\min\{m,r\}},$$

for $\mathrm{Re}(s) > 1$. Since the sum $\sum_{m=M}^{\infty} \sum_p mp^{-m\sigma}$ converges for $\sigma > 1/M$, it follows that the second term of (3.2) can be defined for $\mathrm{Re}(s) > 1/M$ for any M. This gives a meromorphic continuation of $J_r(s,\chi)$ to $\mathrm{Re}(s) > 0$. $\square$

Remark. Using the result in [Da], we see that the imaginary axis is a natural boundary of the function $J_r(s,\chi)$.

4 Asymptotic estimates of group-partitions

To describe the distribution of group-partitions, we recall two lemmas.

Lemma 4.1 (Cf. [Mu] p.26). *For positive integers j and a with $(j,a) = 1$, we have*

$$\frac{1}{\phi(j)} \sum_{\chi \bmod j} \bar{\chi}(a)\chi(b) = \begin{cases} 1 & \text{if} \quad b \equiv a \pmod{j}, \\ 0 & \text{if} \quad b \equiv a \pmod{j}, \end{cases}$$

where χ runs over the distinct Dirichlet characters modulo j, and $\phi(j) = j\prod_{p|j}(1-p^{-1})$ is the Euler function. $\square$

The following lemma is known as the Ikehara-Wiener theorem.

Lemma 4.2 (Cf. [Mu] p.43). *Let $F(s) := \sum_{n=1}^{\infty} b_n n^{-s}$ be the Dirichlet sereis with positive real coefficients and absolutely convergent for $\mathrm{Re}(s) > 1$. Suppose that $F(s)$ can be extended to a meromrphic function in the region $\mathrm{Re}(s) \geq 1$ having no poles except for a simple pole at $s = 1$ with residue $R \geq 0$. Then*

$$\sum_{n \leq x} b_n = Rx + o(x)$$

as $x \to \infty$. $\square$

We now prove the main theorem.

Theorem 4.3. *For $r = 1, 2, 3, \cdots$, or $r = \infty$, we have*

$$\sum_{n \leq x} a_r(n) = \left\{ \prod_{m=2}^{\infty} \zeta(m)^{\min\{m,r\}} \right\} x + o(x) \qquad \text{as } x \to \infty.$$

More precisely, if j and k are positive integers with $(j,k)=1$, then we have

$$\sum_{\substack{n\le x\\ n\equiv k\ (\bmod\ j)}} a_r(n)=\frac{c_r(j)}{\phi(j)}x+o(x)\qquad \text{as } x\to\infty, \tag{4.1}$$

where $\phi(j)$ denotes the Euler function and $c_r(j)$ is a constant which is given by

$$c_r(j):=\prod_{m=1}^{\infty}\prod_{p|j}(1-p^{-m})^{\min\{m,r\}}\times\prod_{m=2}^{\infty}\zeta(m)^{\min\{m,r\}}. \tag{4.2}$$

Proof. For $\mathrm{Re}(s)>1$, we consider the characteristic function:

$$C_r(s,k):=\sum_{\substack{n=0\\ n\equiv k\ (\bmod\ j)}}^{\infty} a_r(n)n^{-s}.$$

From Lemma 4.1, we have

$$\begin{aligned}C_r(s,k)=&\sum_{n=0}^{\infty}\Big\{\frac{1}{\phi(j)}\sum_{\chi}\bar\chi(k)\chi(n)\Big\}a_r(n)n^{-s}=\frac{1}{\phi(j)}\sum_{\chi}\bar\chi(k)J_r(s,\chi)\\ =&\frac{1}{\phi(j)}J_r(s,\chi_0)+\frac{1}{\phi(j)}\sum_{\chi\ne\chi_0}\bar\chi(k)J_r(s,\chi).\end{aligned}$$

We see that $C_r(s,k)$ has an analytic continuation as a meromorphic function to $\mathrm{Re}(s)>0$ with a simple pole at $s=1$. The residue of $C_r(s,k)$ at $s=1$ is given by

$$\begin{aligned}\lim_{s\to1}(s-1)C_r(s,k)&=\frac{1}{\phi(j)}\lim_{s\to1}(s-1)J_r(s,\chi_0)\\ &=\frac{1}{\phi(j)}\prod_{m=1}^{\infty}\prod_{p|j}(1-p^{-m})^{\min\{m,r\}}\times\prod_{m=2}^{\infty}\zeta(m)^{\min\{m,r\}}=\frac{c_r(j)}{\phi(j)}.\end{aligned}$$

Applying Lemma 4.2 to the function $C_r(s,k)$, we obtain the theorem. ☐

For an isomorphic class G of finite Abelian groups, we put $\sigma_r(G)$ by the number of group-partitions of G with just r factors. Then we see that $\sigma_r(G)=\pi_r(G)-\pi_{r-1}(G)$. Let $b_r(n):=\sum_{|G|=n}\sigma_r(G)$. Then, for $r=2,3,4,\cdots$, it is clear that

$$\sum_{\substack{n\le x\\ n\equiv k\ (\bmod\ j)}} b_r(n)=\frac{c_r(j)-c_{r-1}(j)}{\phi(j)}x+o(x)\qquad\text{as } x\to\infty$$

where $c_r(j)$ is a constant given by (4.2) .

Remark. The error term of (1.1) are studied by several authors (Cf. [ES],[Li] for $j = 1$, [Iv],[Sa] for $j \in \mathbb{Z}_{\geq 1}$). Estimations of error terms in (4.1) will be studied in somewhere else.

References

[An] G.E. Andrews, *The theory of partitions.* Encyclopedia of Mathematics and its Applications **2** (1976).

[AS] J.Andersson and J.Snellman, *On the number of plane partitions and non isomorphic subgroup towers of abelian groups.* arXiv:math.NT/0607698, (2006).

[Da] G.Dahlquist, *On the analytic continuation of Eulerian products.* Ark. Mat. **1**, (1952), 533-554.

[ES] P.Erdös, and G.Szekeres, *Über die Anzahl der Abelscher Gruppen gegebener Ordnung und über ein Verwandtes zahlentheoretisches Problem.* Acta Sci. Math. (Szeged) **7** (1935), 97-102.

[Iv] A. Ivić, *An asymptotic formula involving the enumerating function of finite abelian groups.* Univ. Beograd. Publ. Elektrotehn. Fak. Ser. Mat. **3** (1992), 61-66.

[Li] H.Q. Liu, *On the number of abelian groups of a given order.* Acta Arith. **59** (1991), 261-297.

[Mu] M.R. Murty, *Problems in analytic number theory.* Graduate Texts in Mathematics **206** (2001).

[Sa] I.I. Pyateckiĭ-Šapiro, *On an asymptotic formula for the number of Abelian groups whose order does not exceed n.(Russian)* Mat. Sbornik N.S. **26(68)** (1950), 479-486.

Graduate School of Mathematics, Kyushu University
6-10-1, Hakozaki Fukuoka
812-8581, Japan
E-mail momo@math.kyushu-u.ac.jp